


Non-Hispanic Black or African American Health Disparities

Compared to Non-Hispanic Whites In Maryland


Non-Hispanic Blacks or African Americans in Maryland are often more likely than their Non-Hispanic White counterparts to suffer from, and die from, chronic and infectious diseases. These are some of the most important health disparities that afflict Non-Hispanic Blacks or African Americans in Maryland compared to Non-Hispanic Whites.

NON-HISPANIC BLACKS OR AFRICAN AMERICAN HEALTH DISPARITIES: ADULTS



NON-HISPANIC BLACKS OR AFRICAN AMERICAN HEALTH DISPARITIES: CHILDREN

Compared to Non-Hispanic White Children, Non-Hispanic Black or African-American children are more likely to suffer from the following:



*(2006-2010 Age-adjusted mortality rate) Data from CDC Wonder, <http://wonder.cdc.gov>
†(2006-2010 Prevalence) Data from Maryland BRFSS, <http://www.marylandbrfss.org>. Results have been averaged over the following age strata: 18-44 years, 45-64 years, and 65+ years, for the Adult BRFSS metrics.
§ Maryland HIV/AIDS Epidemiological Profile Fourth Quarter 2012, <http://phpa.dhmh.maryland.gov/OIDEOR/CHSE/Shared%20Documents/MarylandHIVepi2013.pdf>